

Review Article

## The Role of Polyphenol Compounds in the Treatment of Cancer Cells

Abu Hazafa<sup>3</sup>, Khalil-Ur-Rehman<sup>1\*</sup>, Nazish Jahan<sup>2</sup> and Zara Jabeen<sup>3</sup><sup>1</sup>Department of Biochemistry, Faculty of Sciences, University of Agriculture, Faisalabad, Pakistan<sup>2</sup>Department of Chemistry, Faculty of Sciences, University of Agriculture, Faisalabad, Pakistan<sup>3</sup>Department of Biochemistry, Faculty of Sciences, University of Agriculture, Faisalabad, Pakistan

\*Corresponding author: Prof. Dr. Khalil-Ur-Rehman, Department of Biochemistry, Faculty of Sciences, University of Agriculture, Faisalabad 38000, Pakistan, Phone: +92-41-5508572, Fax: +92-41-9200764; E-mail: khaliluaf@yahoo.com

Received Date: 01-03-2019

Accepted Date: 12-07-2019

Published Date: 18-08-2019

Copyright: © 2019 Prof. Dr. Khalil-Ur-Rehman

### Abstract

Cancer remains a second leading cause of deaths and major public health problem. It occurs due to extensive DNA damage caused by ultraviolet radiations, ionizing radiations, environmental agents, therapeutic agents, etc. Among all cancers, the most frequently diagnosed cancers are lung (12.7%), breast (10.9%), colorectal (9.7%), and gastric cancer (7.81%). Natural compounds are most favorable against cancer on the count of their anti-cancerous ability, easy to avail and efficient. Among natural compounds, polyphenols (flavonoids, catechin, hesperetin, flavones, quercetin, phenolic acids, ellagic acid, lignans, stilbenes, etc.) represent a large and diverse group used in the prevention and treatment of cancer. Natural polyphenols are derived from different plant sources and from various medicinal plants including *Petroselinum crispum*, *Apium graveolens*, *Flemingia vestita*, *Phyllanthus emblica*, etc. Natural polyphenols possess antioxidant, anti-inflammation, as well as anti-cancerous activities through multiple pathways, they induce apoptosis in breast, colorectal and prostate cancers, lower the nucleoside diphosphate kinase-B activity in lung, bladder and colon cancers, inhibit cell-proliferation and cell cycle arrest by suppressing the NF-kB pathway in various cancers, etc. The current review summarized the anticancer activities of natural polyphenols and their mechanisms of action.

### Abbreviations

Akt pathway: A Serine/Threonine-Protein Kinase pathway; COX-2 :Cyclooxygenase-2; HaCaT :Cultured Human Keratinocyte; HDAC:Histone Deacetylase; MAPK pathway: Mitogen-Activated Protein Kinases pathway (Oestrogen Receptor); NF-kB : *Nuclear factor kappa-light-chain-enhancer of activated B cell*; PARP :Poly ADP Ribose Polymerase; ROS :Reactive oxygen Species; STAT-1:Signal Transducer and Activator of Transcription 1.

**Keywords:** Cancer, Flavonoids; *Phyllanthus emblica*; Lignans; Stilbenes; *Yucca periculosa*


## Introduction

Cancer remains a moment driving reason for death all around the globe after heart diseases [1]. Cancer is an uncontrolled division of ordinary cells in the body. Old cells do not demolish and continue to grow in an uncontrolled way, forming new abnormal cells. These strange cells form a mass of tissue called tumor. Cancer may categorize into five major classes namely as carcinoma, leukemia, sarcoma, lymphomas, and central nervous system cancer [2,3]. The most physiological and biochemical reasons of cancer are ionized and ultraviolet radiations, viral infections (e.g., human papilloma virus HPV cause cervical growth), smoking, parasites (e.g., schistosomiasis cause bladder cancer), contamination of meal or beverages (e.g., liver cancer start by aflatoxins), and bacterial infections (e.g., gastric cancer cause by *Helicobacter pylori*) [4]. Among all cancers, the most frequently diagnosed cancers are lung (12.7%), breast (10.9%), colorectal (9.7%), and gastric cancer (7.81%) [5]. In 2012, approximately 8.2 million people died due to cancer globally, which were associated with liver, breast, cervical, lung, prostate and gastric cancer [6].

Pakistan is the seventh-most crowded nation on the planet, like other developing nations, Pakistan face a double burden of diseases with an expanded rate of cancer. Due to weak database of health system, there are no comprehensive registries available regarding many diseases included cancer [7]. The breast cancer ratio in Pakistan is higher than other Asian countries. In the recent years, the rising rates of head and neck cancer (HNC) are observed and positioned as second most predominant cancer after lung carcinoma in males and breast carcinoma in females in Pakistan [8]. This situation forces the scientists to study about its prevention and treatment. Within the two decades, many laboratories start working on natural polyphenols and their anticancer effects on human being. Polyphenols are potential component for the discovery of anticancer drugs. Polyphenols known as those compounds that have an aromatic ring with at least two or more hydroxyl substituents, including functional derivatives (esters, methyl ethers, glycosides etc.), these compounds are isolated from different kind of fruits (e.g., apples, berries, grapes, oranges, etc.), vegetables (e.g.,

yellow onions, parsley, red peppers, etc.), beverages (e.g., tea, red wine, coffee, etc.) and edible plants. [9-11]. These compounds are water soluble [12] and can be classified into five major classes based on their chemical structure namely as flavonoids, phenolic acids, lignans, stilbenes and other polyphenols. Among of all flavonoids and phenolic acids are the richest [10]. Polyphenol mixes are essential constituents of one the greatest and broadly dispersed group of secondary metabolites in plants [13].

Natural dietary polyphenol compounds may induce their anticancer effects through a variety of mechanisms including the modulation of cell cycle signaling, removal of anti-cancerous agents, antioxidant enzyme activity, apoptosis, and cell cycle arrest (**Figure 1**) [14]. Natural polyphenols modulate the Nrf2 and NF- $\kappa$ B cells [15], an *in vitro* study, shows that polyphenols can significantly affect the MAPK and PI3K cells, which show their participation in cancer cell proliferation [16]. Natural polyphenols, for example, apigenin, resveratrol, genistein, luteolin, and quercetin induced apoptosis in different malignancy cells [17]. Oxidative stress and DNA damage are the most common signals which activate the mitochondrial apoptotic pathway, can rupture the membrane of mitochondria and release the cytochrome C [16].



**Figure 1:** Mechanisms of Natural Polyphenols.

## Prevalence of Cancer

Globally, lung cancer accounts 13% of all malignancies analyzed and 27% of all tumors' death [18]. Prostate cancer is most ordinarily analyzed cancer and the second driving reason for death among men, during 2015, 27,540 people died due to prostate cancer in United State. Breast cancer is the second leading cause of deaths in women after lung cancer with the estimation of 231,840 new cases and 40,730 deaths in 2015. Colorectal cancer is the third most leading cause of death with the estimation of 93,090 and 39,610 cases of colon and rectal cancer respectively, and 49,700 deaths in 2015. The incidence of bladder cancer is almost four times higher among men than the women. Just in 2015, 74,000 new cases anticipated that would be analyzed, and 16,000 deaths occur because of bladder cancer [18, 19].

Epidemiological studies of polyphenol compounds are variable, especially when we consider the results of prospective cohort studies (**Table 1**). Zamora-Ros et al. [20] uncovered that the intake of total flavonoids with lignans significantly reduce the thread of colorectal cancer in Spain. Similarly, a study in Montreal, Canada revealed that the high dietary intake of total flavonoids has inverse association with lung cancer threat [21]. A population-based study in America recommended that the high consumption of antho-

cyanidin (not total flavonoids) may decrease up to 57% of esophageal and gastric cancer [22]. The cohort study of European Prospective Investigation into Cancer and Nutrition (EPIC) revealed that the hepatocellular carcinoma (HCC) risk can be reduced with the intake of flavanols (not total flavonoids), they also observed that total dietary flavonoids are very significant in the reduction of Gastric cancer (GC) risk in women [23,24]. A Cohort study in Netherlands observed that the intake of flavonoids (catechin, epicatechin, kaempferol, myricetin and black tea) significantly reduce the threat of prostate cancer at advanced stage [25]. Badar et al. [26] reported that in Lahore (Pakistan), 15,840 new cancer cases were diagnosed, and 5,134 deaths occurred during 2010-2012, the ratio of male and female patients was 43% and 57% respectively. Akhtar et al. [27] reported that the head and neck (HN) cancer is approximately 8-10% of all cancers in South Asia. According to estimation, the ratio of HN cancer was about 18.74% of all new cancer cases in Pakistan during 2004-2014. Daniyal et al. [28] studied that the stomach cancer risk in Pakistan is low as compared to other countries like US, Japan, etc. Epidemiological data and meta-analysis hardly suggested that when we take diet rich plant polyphenols for long term, they significantly developed our immunity against cancer [29].

**Table 1:** Dietary Natural Polyphenols Consumption and Cancers risk.

Types of Cancer	Polyphenols	Study Type	Risk Estimation (95% CI)	References
Lung cancer	Flavonoids	case-control study	0.63 (0.47-0.85)	[21]
Prostate cancer	Flavonoids	cohort study	Total catechin 0.73 (0.57-0.95), Kaempferol 0.78 (0.61-1.00), Epicatechin 0.74 (0.57-0.95)	[25]
Breast cancer	Flavonoids	meta-analysis	Flavonols 0.88 (0.88-0.98), Flavones 0.83 (0.76-0.92)	[30]
	Flavanols	cohort study	0.81 (0.67-0.97)	[31]
	Isoflavones	meta-analysis	0.68 (0.52-0.89)	[32]
Colorectal cancer	Flavonoids and Lignans	case-control study	Total flavonoids 0.59 (0.38-0.99); Lignans 0.59 (0.34-0.99)	[20]
	Isoflavones	meta-analysis	0.76 (0.59-0.99)	[33]
HCC	Flavanols	cohort study	0.62 (0.33-0.99)	[23]
Gastric cancer	Flavonoids	case-control study	0.33 (0.15-0.73)	[34]
	Flavonoids	case-control study	No significant association	[22]

## Treatment of cancer by natural compounds

Different types of therapies are available to treat the cancer, but they can induce the toxicity and side effects due to their specificity at a single target [35]. Cancer is a multistage disease, so those compounds are required for the prevention and treatment of cancer, which can target the multiple molecular and biochemical pathways without creating any poisonous quality and reactions to ordinary cells [36]. Cancer chemoprevention is the major field in the prevention of carcinomas using different pharmaceutical, synthetic and natural compounds to completely retard, reverse or inhibit the process of carcinomas. Natural compounds are most favored for the treatment of carcinogenesis on the count of their anti-cancerous ability, ease of availability, potential to overcome the resistance, safety, and efficiency [37]. The Plant kingdom are best sources to anti-cancer drugs. Around thirty different anti-cancerous natural mixes have been separated from plants and more than 3,000 species of plants have been accounted in the treatment of cancer and clinical trials so far. Today about

25% of all traditions bear one or more components from plants [38]. Numerous secondary metabolites which are expelled from plant sources represent the potential novel agents by inducing the apoptosis and cell cycle arrest [13]. Among all, natural polyphenol compounds (flavonoids, phenolic acids, lignans, and stilbenes) represent a large and diverse group in the prevention and treatment of cancer [39].

Polyphenols are natural, but synthetic and organic compounds, they always have heteroatom substitutes other than hydroxyl groups. Natural Polyphenol compounds classified into five groups such as flavonoids, phenolic acids, lignans, stilbenes and other polyphenols (Table 2) [10]. These compounds are found almost in all families of plant and can also be isolated from an aquatic submerged plant *Myriophyllum spicatum* that involved in allopathic interactions [40]. Polyphenols also have ability to bind with the proteins and form soluble and insoluble protein-polyphenol complexes [41].

**Table 2:** Classification of Natural Polyphenols.

Classification	Represented members	Some common food sources
Anthocyanidins	Cyanidin, Delphinidin, Malvidin, Pelargonidin, Peonidin, Petunidin	red, blue, and purple berries; red and purple grapes; red wine
Flavanols	Monomers: Catechin, Epicatechin, Epigallocatechin Epicatechin gallate Dimers and Polymers: Theaflavins, Thearubigins, Proanthocyanidins	Catechins: Teas (particularly green and white), chocolate, grapes, berries, apples Theaflavions and Thearubigins: Teas (particularly black) Proanthocyanidins: Apples, chocolate, red wine, red grapes
Flavanones	Hesperetin, Naringenin, Eriodictyol	Citrus fruits and juices, e.g., oranges, grapefruits, lemons
Flavonols	Quercetin, Kaempferol, Myricetin, Isorhamnetin	Widely distributed in yellow onions, scallions, kale and tea
Flavones	Apigenin, Luteolin	Parsley, thyme, celery, red peppers
Isoflavones	Daidzein, Genistein, Glycitein	Soybeans, soy foods, legumes
Phenolic acids	Hydroxybenzoic Acid	Ellagic acid, Gallic acid Grapes, pomegranates and chocolate
	Hydroxycinnamic Acid	Ferulic acid, Chlorogeni acid Coffee, cereal grains
Lignans	Sesamin, Secoisolariciresinol, Diglucoside	Flaxseeds, sesame
Stilbenes	Resveratrol, Pterostilbene, Piceatannol	Grapes, berries, red wine

This information is taken from the references [9,12]


### Typical natural polyphenols and their anticancer activities

Polyphenols are found in foodstuffs and are known to possess anticancer properties; these compounds are derived from secondary metabolites for the protection against predations, ultraviolet radiations, plant pigmentation and reproduction. These mixes are ubiquitous in vegetables, fruits, and drinks [42,43]. Following we will summarize all the classes of natural polyphenols and their anticancer effects:

#### Flavonoids

Flavonoid is a wide class of polyphenolic compounds; the main sources of flavonoids are fruits, vegeta-

bles, seeds, flowers and beverages, different type of foods have enough quantity of anti-cancerous flavonoid (Table 3). The chemical structure of natural flavonoid is diphenylpropane (C6-C3-C6), which contain two aromatic rings at both corner and a three-carbon ring at the center, which forming an oxygenated heterocyclic (Figure 2) [44]. The best-known biological effects of natural flavonoids are prevention against cancer, inhibition of bone resorption, cardio-protective and hormonal action. In addition, they are also more effective against antiviral and antibacterial properties [45,46].



**Figure 2:** The representative structures of Flavonoid.

**Table 3:** Foods that contain high amounts of natural Flavonoids.

Description	Class	Subclasses	mg/100g	References
<b>Onions</b>	Flavonols	Kaempferol	0.7	[48]
		Myricetin	2.16	
	Flavones	Apigenin	0.24	
		Luteolin	0.16	
<b>Grape fruit, raw</b>	Flavanones	Naringenin	53	[49]
		Hesperetin	1.5	
	Flavonols	Quercetin	0.5	
<b>Soybeans, mature seeds</b>	Isoflavones	Genistein	80.99	[50]
		Daidzein	62.07	
<b>Cacao beans</b>	Flavan-3-ols	(-)-Epicatechin	99.18	[51]
		(+)-Catechin	88.45	
<b>Spices, parsley, dried (<i>Petroselinum crispum</i>)</b>	Flavones	Apigenin	4503.5	[52]
		Luteolin	19.75	
<b>Green tea, brewed, decaffeinated</b>	Flavan-3-ol	(-)-Epigallocatechin 3-gallate	26.05	[53]
	Flavonols	Quercetin	2.77	
		Kaempferol	1.00	
<b>Black tea, brewed, (prepared with tap water)</b>	Flavan-3-ols	(-)-Epigallocatechin 3-gallate	9.36	[54]
		(+)-Catechin	1.51	
	Flavonols	Quercetin	2.19	
		Myricetin	0.45	

The affinity of flavonoids for cellular membrane depends on the glycosylation and number of hydroxyl groups, substitutive groups also have great impact on their structure, fluidity and permeability. In the metabolism and conjugation of total flavonoids inside the human, the gastrointestinal tract and the colonic micro-flora play an important role and help the total flavonoids into the systematic circulation and liver [47]. Flavonoids can also bind with the ATP-binding sites of assortments of proteins including mitochondrial ATPase, protein kinase A, protein kinase C, calcium plasma membrane ATPase and topoisomerase [44].

### Flavanols

Flavanols represent a restricted group of flavonoids, which contain the 2-phenylchromanol skeleton, and can be divided into three main categories monomers, oligomers and polymers. Flavanols are commonly found in fruits, vegetables or some beverages, and shows various biological activities that render them useful for health [55].

### Green tea catechins as anticancer drug

Green tea catechins contain (-)-epigallocatechin (EGC), (-)-epicatechin gallate (ECG), and (-)-epigallocatechin gallate (EGCG) [56]. Tea catechins are powerful inhibitors of cell proliferation by inhibiting the acceleration of cells expressing cancer-specific genes, they are also potent inhibitors of cancer metastasis by lowering the nucleoside diphosphate kinase-B activity, due to these activities they have been used for the treatment of lung, bladder, colon, ocular, and cervical cancers which caused by human papilloma virus [57]. Epigallocatechin gallate (EGCG), is one of the most extensively investigated components of green tea, which have numerous advantageous impacts to lessen the tumor and cardiovascular diseases. EGCG founds in dried leaves of green tea (7,380 mg/100g), white tea (4,245 mg/100g), and also found in black tea in very mute quantity (936 mg/100g) [58]. Herbitual green tea intake provides some relief against breast cancer and chemo-prevention in prostate [59].

A study showed that in vitro the EGCG treatment could suppress the nicotine-induced migration and invasion of A549 lung cancer cells [60]. EGCG virtually suppressed the invasion and migration of CL1-5 lung cancer cells by suppressing the matrix metalloproteinase-2 (MMP-2) expression at concentration of 5-20  $\mu$ M [61]. Banerjee [62] suggested that EGCG and theaflavins (TF) together were assessed for their chemo-preventive potential, these two compounds could activate the caspase-3, caspase-7, and Cox-2 expression, as a result of that, and they lowered the incidence of pre-invasive lung cancer.

Sen et al. [63]; Khan and Mukhtar [64] reported that the Mg<sup>2+</sup> plays an important role in increasing the inhibitory effect of EGCG on human DNMT activity. In the context of human cell lines, EGCG (20  $\mu$ mol/L) inhibits the activity of DNMT in breast (MDA-MB-231 and MCF-7), prostate (PC-3), esophageal (KYSE-150), and colon (HT-29) cancer cells. Chuang et al. [65]; Gilbert and Liu [66] uncovered that the treatment of breast (MDA-MB-231 and MCF-7) cancer cells is performed by the down-regulation of human telomerase reverse transcriptase (hTERT) expression which lead to reduce the telomerase level.

### **Anthocyanidins**

Anthocyanidins are excessively present in the plant kingdom. Red wine, purple cabbage, berries, and grapes are the main sources of anthocyanidins, it is classified into six further classes such as cyanidin, delphinidin, pelargonidin, malvidin, petunidin, and peonidin [67]. It is reported that anthocyanidins can reduce risk of cancer, arthritis, and cardiovascular diseases due to its anti-mutagenesis, antioxidant, and anti-inflammatory activities. A recent study uncovered that the ortho-dihydroxyphenyl structure on the ring number B is responsible to inhibit the growth of tumor and metastasis [68].

### **Delphinidin as anticancer drug**

Among of all anthocyanidins, delphinidin have the highest anticancer activity. Delphinidin suppress the NF- $\kappa$ B pathway to induce apoptosis and cell cycle arrest in several types of cancer [69]. Kausar et al. [70] indicated that the anthocyanidins might show the better role in the com-

bination with other compound than the single one in the treatment of cancer. A study uncovered that the anthocyanidins (peonidin-3-glucoside, cyaniding-3-glucoside and cyanidin-3-O-sambubioside) that are expelled from black rice and the fruit *acanthopanax sessiliflorus*, could induce the cell-proliferation, apoptosis, and tumor growth of HER2 cells, and inhibit the angiogenesis and invasion in the positive breast cancer [71,72]. Similarly, cyanidin and delphinidin shows the oxidative stress-based cytotoxicity to colorectal cancer cells [72].

### **Flavonols**

The main sources of flavonols (3-hydroxy-2-phenylchromen-4-one) are onions, kale, teas, berries and apples but they are usually present at low concentration, it is further classified into different subgroups namely as quercetin, kaempferols, myricetin and isorhamnetin etc. [73]. Flavonols show their anti-cancer activities through various mechanisms including the antioxidant activity, interactions with proteins, and inhibit the enzymes (CYP2C9 and CYP3A4) that regulates the cancer cells [74].

### **Quercetin as anticancer drug**

Quercetin (2-phenyl-4H-1-benzopyran-4-one, 2-phenylchromone) is a bioactive and predominant flavonol, its chemical structure is responsible for its anti-oxidant activities, the hydroxyl groups that attached at heterocyclic ring B at carbon 3 and 5 are responsible for its anti-proliferative potential and also give good sensitivity against the free-radicals [36,75]. A study indicated that the daily intake of quercetin in the United States, Europe, and Asia is ranging from 4 to 68 mg per individual [76]. The metabolite of quercetin (quercetin-3-O-glucuronide), have the significant role in inhibition the noradrenaline, which bind with 2-adrenergic by suppressing the DNA damage, that is induced during the treatment of human breast cancer cells (MCF-10A) by noradrenaline and 4-hydroxyestradiol [77].

The best possible known mechanism of quercetin is not completely found yet, but the least investigated anti-cancer outcomes of quercetin is highlighted as it could repress the sodium-dependent glucose transporter-1 (SGLT-1), block the cell cycle at several phases (G2/M or G1), and

can go about as a competitive inhibitor of efflux pumps. In short, it promotes the apoptosis [36].

### Flavanones

Flavanones (2-phenylchroman-4-ones) are most abundant in the peel and the juices of citrus fruits, and additionally ordered into various subclasses namely as naringenin, hesperetin, eriodictyol, etc. [78]. Flavanones possess many pharmacological properties including antibacterial, antioxidant, anti-fungal and antiVSMCs vegetation activities, and have been reported as a potent anticancer agent [79].

#### Hesperetin as anticancer drug

Hesperetin belong to class flavanones, mainly founds in citrus fruits included oranges and grapefruits. Hesperetin could reduce the DNMT activity in the nuclear extraction of esophageal squamous cell carcinoma KYSE-510, when it incubated for 1.5h with 20-50  $\mu\text{mol/L}$  [74]. Choi [80] recommended that the hesperetin capture the cell cycle at  $G_1$ -stage during the treatment of breast cancer (MCF-7) cells and suppress the activity of cell proliferation in carcinoid cells by expending the Notch-1 expression. Palit et al. [81] uncovered that in the treatment of breast cancer cells, hesperetin (40-200  $\mu\text{M}$ ) promote the ROS and ASK1/JNK activation pathways and induce the growth inhibition through mitochondria-mediated apoptosis.

In cervical malignancy cells hesperetin initiated apoptosis through death receptors and mitochondrial pathways [82]. In the prostate cancer PC-3 cells hesperetin (40-90  $\mu\text{M}$ ) induced the apoptosis, which are mediated by the NF- $\kappa\text{B}$  pathway [83]. Zhang et al. [84] demonstrated that the hesperetin (100-400  $\mu\text{M}$ ) induced the apoptosis (mitochondria-mediated) through promoting the ROS accumulation in the treatment of gastric tumor cells.

### Flavones

Flavones are a class of total flavonoids with the backbone of 2-phenylchromen-4-one (2-phenyl1-benzopyran-4-one), they are usually found in products of the soil like celery, thyme, parsley, etc., and show their solubility toward water and ethanol. Because of their anti-proliferative,

antioxidant, and anti-inflammatory activities, they could diminish the threat of cancer and cardiovascular ailments [78,85].

#### The anticancer activity of apigenin

Apigenins are derived from some medicinal plants such as *Lycopodium clavatum* (club moss), and from some vegetables, for example, *Apium graveolens* (celery) and *Petroselinum crispum* (parsley) [86]. Recently *in vitro* study reported that apigenin have numerous gainful roles in anti-proliferation properties with the enlistment of apoptosis and cell cycle arrest in the cancer cells, and also shown down-regulation of NF- $\kappa\text{B}$ , UV-B-induced ROS, and DNA damage by removing the cyclobutane rings [87].

Apigenin initiated the apoptosis and cell cycle arrest in the treatment of prostate cancer (PC-3 and 22Rv1) cells. The daily dose of apigenin (20-40  $\mu\text{mol/L}$ ) suppressed the activity of HDAC1 and HDAC3 on protein level and mRNA which resulted the hyperacetylation of histone (H3 and H4) during the treatment of prostate cancer [88]. Apigenin possess the protective effect in epithelial cells from lipopolysaccharide (LPS)-induced inflammation. It holds the DNA damage under wraps by reducing the caspase-3 expression levels and ROS production [89]. Apigenin initiates the ATM and H2AX phosphorylation with the down-regulation of cell cycle controlling and DNA repair genes, which incites the apoptosis/DNA damage in the malignancy cells [90]. Kroonen et al. [91] demonstrated that apigenin control the DNA damage by the activation of NF- $\kappa\text{B}$  in malignant glioma cells, inhibition of casein kinase-2, and regulations of cell proliferation. Gates et al. [92] suggested in a population-based examination that ovarian cancer can be overcome by apigenin.

#### Isoflavones

The main products of Isoflavones are peas, legumes, and alfalfa. Genistein and daidzein are the most prominent isoflavones. Isoflavones shown anticancer activities against leukemia, prostate, breast, colon and lung cancer [93].

#### The anticancer activity of genistein


Genistein was first isolated from dyer's broom *Genista tinctoria* plant. The main sources are soybeans, fava

beans, and psoralea. Also founds in many medicinal plants like *Flemingia vestita* and *Flemingia macrophylla* [94]. Genistein indicates chemo-therapeutic and chemo-preventive exercises in numerous tumors, particularly in hormone-dependent cancers. The *in vivo* and *in vitro* studies conformed its antineoplastic effects in prostate, breast, urinary, colon, and skin cancer [95]. Tian et al. [96] reported that in lung cancer treatment genistein (25-75  $\mu\text{M}$ ) could inhibit the cell proliferation and migration of H446 cell by the induction of cycle arrest at  $G_2/M$ -stages and apoptosis. In the prostate malignancy (LNCaP and PC-3) and esophageal squamous cell carcinoma (KYSE-150), genistein repress the DNMT by the reduction of cellular accessibility of SAM with the dose of 20-50  $\mu\text{mol/L}$  [97]. In the breast cancer the conjugates of EGF-genistein could inhibit the EGF-Rc tyrosine kinase and induce the apoptosis, killing approximately 99.99% of clonogenic breast cancer cells [98]. A study revealed that in the treatment (25-100  $\mu\text{M}$ ) of colon cancer cells, genistein exhibit pro-apoptosis and anti-proliferative effects [99].

### Phenolic acids

Phenolic acids are aromatic acidic compounds and are significant constituent of plants which are derived from a variety of fruits, vegetables, beverages, legumes and from other edible plants. Hydroxybenzoic acid (gallic acids, ellagic acids, etc.) and hydroxycinnamic acid (cinnamic acids, caffeic acids, ferulic acids, etc.) are chief subclasses of phenolic acids (**Figure 3**) [100]. Phenolic acids and its an-


alogues namely as Gallic and caffeic acids, display verities of biological activities, in addition to their antioxidant activities, which are related to regulations of carcinogenesis [101].



**Figure 3:** The representative structures of Phenolic acid.

### Hydroxybenzoic acids

Hydroxybenzoic acids are available in couple of edible plant and its concentration is exceptionally factor with the utilization of coffee. The derivatives of Hydroxybenzoic acid (protocatechuic, vanillic and p-hydroxybenzoic acids) form p-coumaric and caffeic acid, which have important anticancer activities [102].




**Figure 4:** The Chemical structures of lignan derivatives.

### Gallic acid as anticancer drug

Gallic acid is widely extracted from plant-based fruits and present in hydrolyzable tannins in minute quantity. It possesses numerous pharmacological (anticancer,

antimicrobial and anti-inflammatory) activities, and induced apoptotic in colon cancer cells (HCT-15) [103]. Gallic acids are segregated from *Phyllanthus emblica* fruit, which demonstrated its anti-proliferative exercises against breast



**Figure 5:** The Chemical structures of Stilbene derivatives.

### Anticancer activity of resveratrol:

Resveratrol (3,4,5-trihydroxy-trans-stilbene) is an important polyphenol, berries, grapes, peanuts and other edible plants are the major products of resveratrol. Resveratrol also expelled from the root of *Polygonum cuspidatum* [120]. Resveratrol possess multiple pharmacological, molecular and biochemical properties against precancerous cells. *In vitro* and *in vivo* studies recommended that resveratrol have properties against a variety of human ailments, including cardio and neuro-protection, invulnerable direction, and cancer chemoprevention [122]. Resveratrol

prompted apoptosis and cell cycle arrest in precancerous or cancer cells by controlling the different signal pathways, including Akt, MAPK, Wnt, etc., without effecting the normal cells [123]. In the treatment of gastric cancer cells, resveratrol (20-50 $\mu$ M) actuated the cell cycle capture at G<sub>1</sub> stage [124]. In the chemo-resistant human colon cancer cells, resveratrol (50-400 $\mu$ M) used for 30 minutes as anti-cancer drugs, which induce apoptosis via CD95 and Wnt pathways [125]. In the breast cancer, resveratrol induced apoptosis in MCF-7 and casp-3 cells [126]. Some cancers and their pathways are enlisted in **Table 4**.

**Table 4:** Resveratrol involved in different pathways against multiple Cancers.

Cancers	Pathways	References
Breast cancer	CD95	[127]
Colon cancer	CD95, Wnt	[128]
Lung cancer	NF- $\kappa$ B, SIRT1	[129]
Prostate cancer	NF- $\kappa$ B	[130]
Lymphocytes	CD95	[123]
Skin cancer	Apoptosis, NF-Kb	[131]

cancer cells. In stomach adenocarcinoma, gallic acids (3.5  $\mu\text{M}$ ) reduce the tumor metastasis by inhibiting the activities of NF- $\kappa\text{B}$  and down-regulation of PI3K/AKT cells [104]. In leukemia, gallic acid showed pro-apoptosis activities by the production of hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) [105]. Zhao and Hu [106] reported that gallic acid (10-40  $\mu\text{g}/\text{ml}$ ) significantly reduce the cell-proliferation, invasion and angiogenesis HTB-35 and HeLa in the treatment of cervical cancer. Forester et al. [107] demonstrated that in the treatment of colon cancer cells (Caco-2) gallic acid induce apoptosis and

cell cycle rest at  $G_0/G_1$  stage by inhibiting the transcriptional factors (STAT-1, NF- $\kappa\text{B}$  and OCT-1).

### Hydroxycinnamic acids

Hydroxycinnamic acids belong to the class aromatic acids and classified into many subgroups such as caffeic, cinnamic, chlorogenic, coumaric, sinapic and ferulic acids. In cancer cells they showed inhibitory cell-proliferation activity [108].

**Table 5:** Natural Polyphenols as Anticancer in the in vivo and in vitro studies.

Polyphenols/ Cancer Type	Study Type	Dose	References
<b>Breast cancer</b>			
EGCG	<i>in vitro</i>	1–40 $\mu\text{M}$	[135]
Hesperetin	<i>in vitro</i>	40–200 $\mu\text{M}$	[81]
Chrysin	<i>in vitro</i>	5–20 $\mu\text{M}$	[136]
Quercetin	<i>in vitro</i>	1–200 $\mu\text{M}$	[137]
Luteolin	<i>in vitro</i>	10–40 $\mu\text{M}$	[94]
Daidzein	<i>in vitro</i>	3–50 $\mu\text{M}$	[138]
Genistein	<i>in vitro</i>	5–20 $\mu\text{M}$	[139]
Genistein	<i>in vivo</i>	$\leq 500$ ppm	[140]
Ellagic acid	<i>in vitro</i>	10–40 $\mu\text{g}/\text{mL}$	[141]
Ellagic acid	<i>in vivo</i>	50–100 mg/kg	[94]
<b>Lung cancer</b>			
Anthocyanidins	<i>in vivo</i>	0.5 mg/mouse	[70]
Xanthohumol	<i>in vitro</i>	14–42 $\mu\text{M}$	[142]
EGCG	<i>in vitro</i>	5–20 $\mu\text{M}$	[61]
Naringenin	<i>in vitro</i>	100 $\mu\text{M}$	[143]
Luteolin	<i>in vivo</i>	10–30 mg/kg	[144]
Genistein	<i>in vitro</i>	25–75 $\mu\text{M}$	[96]
Resveratrol	<i>in vitro</i>	5–50 $\mu\text{M}$	[130]
<b>Prostate cancer</b>			
EGCG	<i>in vivo</i>	1 mg 3 $\times$ /week	[145]

Apigenin	<i>in vivo</i>	20 and 50 µg/ mouse	[146]
Genistein	<i>in vitro</i>	0.5–50 µM	[140]
Ellagic acid	<i>in vitro</i>	10–100 µM	[147]
Sesamin	<i>in vivo</i>	10 mg/kg	[119]
Resveratrol	<i>in vivo</i>	30 mg/kg	[148]
Resveratrol	<i>in vitro</i>	25–100 µM	[149]
<b>Colorectal cancer</b>			
Delphinidin	<i>in vitro</i>	30–240 µM	[150]
EGCG	<i>in vitro</i>	1–50 µM	[151]
Naringenin	<i>in vitro</i>	50–200 µM	[152]
Apigenin	<i>in vivo</i>	50 mg/kg	[153]
Luteolin	<i>in vitro</i>	20–100 µM	[154]
Kaempferol	<i>in vitro</i>	0–60 µM	[155]
Genistein	<i>in vivo</i>	20–80 mg/kg	[99]
Resveratrol	<i>in vitro</i>	25–150 µM	[156]
<b>Cervical cancer</b>			
Hesperetin	<i>in vitro</i>	650 µM	[82]
Genistein	<i>in vitro</i>	100 µM	[157]
Quercetin	<i>in vitro</i>	110.38 µM	[158]
Resveratrol	<i>in vitro</i>	150–250 µM	[159]
<b>Liver cancer</b>			
Xanthohumol	<i>in vitro</i>	5–40 µM	[160]
Daidzein	<i>in vitro</i>	200–600 µM	[161]
Sesamin	<i>in vitro</i>	25–125 µM	[118]
Ellagic acid	<i>in vivo</i>	30 mg/kg	[162]
Resveratrol	<i>in vitro</i>	25–100 µM	[156]
Resveratrol	<i>in vivo</i>	20 mg/kg	[163]
<b>Gastric Cancer</b>			
EGCG	<i>in vitro</i>	20–100 µM	[164]
Hesperetin	<i>in vivo</i>	20–40 mg/kg	[84]
Kaempferol	<i>in vivo</i>	20 mg/kg	[155]
Apigenin	<i>in vitro</i>	20 µg/mL	[165]
Gallic acid	<i>in vivo</i>	0.25% and 0.5% in water	[107]

**Table 6:** Mechanism of actions of Natural Plant derived Polyphenols.

Mechanism of Actions	Phenolic compounds	References
<b>Antiangiogenesis and antimutagenic properties</b>		
Inhibit cell proliferation, inhibit oncogene expression, induce tumor suppressor gene expression, and DNA binding prevention.	Favonoids (apigenin, daidzein, luteolin, kaempferol, myricetin, quercetin, hesperetin, EGCG, and genistein), Phenolic acids (chlorogenic acid and hydroxytyrosol), stilbenes (resveratrol), and lignans.	[166,167]
<b>Induction of cell-cycle arrest and apoptosis</b>		
Inhibit cell cycle at different phases: S/G <sub>2</sub> , G <sub>1</sub> , and S phases, direct or indirect effect on cell cycle arrest, involved in p53 and induce apoptosis.	Flavonoids (quercetin and EGCG), Phenolic acids (ferulic acid, caffeic acid, ellagic acid, and hydroxytyrosol), Stilbenes (resveratrol), and lignans (sesamin).	[166,168]
<b>Inhibition of signal transduction pathways</b>		
NF-κB and AP-1 signaling pathway, Nrf-KEAP1 (Kelch-like ECH-associated protein 1) complex signaling pathways, the Wnt or β-catenin signaling pathway, growth-factor receptor-mediated pathways, and MAPK signaling pathway.	Flavonoids (apigenin, genistein, quercetin, and EGCG), phenolic acids (caffeic acid and gingerol), lignans, coumarins, and stilbenes (resveratrol).	[168,169]
<b>Antiviral, antibacterial, and antifungal effects</b>		
Inhibit liver cancer and down-regulate HIV expression	Flavonoids, phenolic acids, stilbenes (resveratrol), and lignans.	[167]
<b>Enzyme inhibition</b>		
COX-2, iNOS, XO, signal transduction Enzymes such as PKC and PTK, phase I enzyme (block activation of carcinogens), and DNA methyltransferases.	Flavonoids (apigenin, luteolin, quercetin, and EGCG), Phenolic acids (chlorogenic acid, caffeic acid, ellagic acid), stilbenes (resveratrol), and lignans.	[166,167]

---

## Conclusion and Future Development

Cancer is a mind-boggling process which includes in the multiplication of cells, hindered cells passing, and worldly changes in the physiology of cells. Cancer remains a moment driving reason for death comprehensively. Malignancy happens because of dysfunction of numerous coding genes proteins including development factor receptors, hostile to apoptotic proteins, and so on. Lung, breast, colorectal, and stomach cancers are most prominent cancer among all. Pakistan has the most elevated breast cancer proportion among Asian nations, and most normal tumors in Pakistan are lung, breast, head and neck cancers, and lymphoma. Natural compounds are most supported for the treatment of carcinogen because of their security, proficiency, straightforwardness to accessibility, and hostile to malignant capacity. Phytochemicals shown a significant role against various malignancies, but among all, natural polyphenol compounds (flavonoids, phenolic acids, lignans, and stilbenes) represent a broad and assorted group in the anticipation and treatment of cancer. Natural polyphenols have their anticancer exercises through various pathways, including cancer prevention agent enzymatic movement, cell cycle arrest (S/G<sub>2</sub>, G<sub>1</sub>, S, and G<sub>2</sub> stages), by regulating the Nrf2 and NF-κB cells, inhibit the cell-proliferation, prompt apoptosis, MAPK flagging pathway, and so on. The anticancer impacts of polyphenol changed with the doses, cancer type, and cell lines.

The clinical trials are limited about the natural polyphenol compounds. In the future, more epidemiological investigations employing biomarkers of polyphenols are needed to assess the effect of dietary polyphenols on cancer hazard. It is recommended that the synthesis of well-targeted and well-designed phenolic compounds, can lead to the development of clinically beneficial drugs with efficiency and selectivity. In addition, various researches are required to investigate that which signals, and mechanisms are responsible for the regulation of apoptosis and cell cycle death after the treatment by using natural polyphenols.

## Acknowledgements

The author is extremely grateful to ALMIGHTY ALLAH, who in spite of numerous difficulties and acute frustrations enabled me to complete the review article. The author is also thankful to Prof. Dr. Khalil-ur-Rehman and Dr. Zara Jabeen for reviewing the article and providing the helpful comments. This article is supported by Department of Biochemistry from University of Agriculture, Faisalabad, Pakistan.

## References

1. Millimouno FM, Dong J, Yang L, et al. Targeting apoptosis pathways in cancer and perspectives with natural compounds from mother nature. *Cancer Prev Res* 2014; 12.
  2. Esteller M. Epigenetics in cancer. *N Engl J Med* 2008; 358: 1148-1159.
  3. Parker JS, Mullins M, Cheang MC, et al. Supervised risk predictor of breast cancer based on intrinsic subtypes. *J Clin Oncol* 2009; 27: 1160.
  4. Block KI, Gyllenhaal C, Lowe L, et al. Designing a broad-spectrum integrative approach for cancer prevention and treatment. In, *Seminars in cancer biology: Elsevier* 2015; S276-S304.
  5. Bray F, Ren JS, Masuyer E, et al. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *Int J Cancer* 2013; 132: 1133-1145.
  6. WHO Cancer 2016; Available online: <http://www.who.int/mediacentre/factsheets/fs297/en/>.
  7. Hanif M, Zaidi P, Kamal S, et al. Institution-based cancer incidence in a local population in Pakistan: nine years data analysis. *Asian Pac J Cancer Prev* 2009; 10: 227-230.
  8. Nosheen M, Ishrat M, Malik F, et al. Association of GSTM1 and GSTT1 gene deletions with risk of head and neck cancer in Pakistan: a case control study. *Asian Pac J Cancer Prev* 2010; 11: 881-885.
-

- 
9. Manach C, Scalbert A, Morand C, et al. Polyphenols: food sources and bioavailability. *Am J Clin Nutr* 2004; 79: 727-747.
  10. Neveu V, Perez-Jiménez J, Vos F, et al. Phenol-Explorer: an online comprehensive database on polyphenol contents in foods. *Database* 2010; 2010: 024.
  11. D'Archivio M, Filesì C, Vari R, et al. Bioavailability of the polyphenols: status and controversies. *Int J Mol Sci* 2010; 11: 1321-1342.
  12. Quideau S, Deffieux D, Douat-Casassus C, et al. Plant polyphenols: chemical properties, biological activities, and synthesis. *Angew Chem Int Ed Engl* 2011; 50: 586-621.
  13. Marín L, Miguélez EM, Villar CJ, et al. Bioavailability of dietary polyphenols and gut microbiota metabolism: antimicrobial properties. *Biomed Res Int* 2015; 15: 1-18.
  14. Ramos S. Cancer chemoprevention and chemotherapy: dietary polyphenols and signalling pathways. *Mol Nutr Food Res* 2008; 52: 507-526.
  15. Gopalakrishnan A, Kong A-NT. Anticarcinogenesis by dietary phytochemicals: cytoprotection by Nrf2 in normal cells and cytotoxicity by modulation of transcription factors NF- $\kappa$ B and AP-1 in abnormal cancer cells. *Food Chem Toxicol* 2008; 46: 1257-1270.
  16. Dhillon AS, Hagan S, Rath O, et al. MAP kinase signalling pathways in cancer. *Oncogene* 2007; 26: 3279.
  17. Vauzour D, Rodriguez-Mateos A, Corona G, et al. Polyphenols and human health: prevention of disease and mechanisms of action. *Nutrients* 2010; 2: 1106-1131.
  18. Torre LA, Bray F, Siegel RL, et al. Global cancer statistics 2012. *CA Cancer J Clin* 2015; 65: 87-108.
  19. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin* 2015; 65: 5-29.
  20. Zamora-Ros R, Not C, Guinó E, et al. Association between habitual dietary flavonoid and lignan intake and colorectal cancer in a Spanish case-control study (the Bellvitge Colorectal Cancer Study). *Cancer Causes Control* 2013; 24: 549-557.
  21. Christensen KY, Naidu A, Parent MÉ, et al. The risk of lung cancer related to dietary intake of flavonoids. *Nutr Cancer* 2012; 64: 964-974.
  22. Petrick J, Steck S, Bradshaw P, et al. Dietary intake of flavonoids and oesophageal and gastric cancer: incidence and survival in the United States of America (USA). *Br J Cancer* 2015; 112: 1291.
  23. Zamora-Ros R, Fedirko V, Trichopoulou A, et al. Dietary flavonoid, lignan and antioxidant capacity and risk of hepatocellular carcinoma in the European prospective investigation into cancer and nutrition study. *Int J Cancer* 2013; 133: 2429-2443.
  24. Zamora-Ros R, Agudo A, Luján-Barroso L, et al. Dietary flavonoid and lignan intake and gastric adenocarcinoma risk in the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Am J Clin Nutr* 2012; 96: 1398-1408.
  25. Geybels MS, Verhage BA, Arts IC, et al. Dietary flavonoid intake, black tea consumption, and risk of overall and advanced stage prostate cancer. *Am J Epidemiol* 2013; 177: 1388-1398.
  26. Badar F, Mahmood S. Epidemiology of cancers in Lahore, Pakistan, among children, adolescents and adults, 2010-2012: a cross-sectional study part 2. *BMJ Open* 2017; 7: e016559.
  27. Akhtar A, Hussain I, Talha M, et al. Prevalence and diagnostic of head and neck cancer in Pakistan. *Pak J Pharm Sci* 2016; 29: 1839-1846.
  28. Daniyal M, Ahmad S, Ahmad M, et al. Risk factors and epidemiology of gastric cancer in Pakistan. *Asian Pac J Cancer Prev* 2015; 16: 4821-4824.
-

- 
29. Pandey K, Rizvi S. Plant polyphenols as dietary antioxidants in human health and disease. *Oxid Med Cell Longev* 2009; 2: 270-278.
  30. Hui C, Qi X, Qianyong Z, et al. Flavonoids, flavonoid subclasses and breast cancer risk: a meta-analysis of epidemiologic studies. *PLoS One* 2013; 8: e54318.
  31. Wang Y, Gapstur SM, Gaudet MM, et al. Evidence for an Association of Dietary Flavonoid Intake with Breast Cancer Risk by Estrogen Receptor Status is Limited-3. *J Nutr* 2014; 144: 1603-1611.
  32. Xi Q, Chen M-L, Qin Y, et al. Isoflavone consumption and risk of breast cancer: a dose-response meta-analysis of observational studies. *Asia Pac J Clin Nutr* 2013; 22: 118-127.
  33. Tse G, Eslick GD. Soy and isoflavone consumption and risk of gastrointestinal cancer: a systematic review and meta-analysis. *Eur J Nutr* 2016; 55: 63-73.
  34. Woo HD, Lee J, Choi IJ, et al. Dietary flavonoids and gastric cancer risk in a Korean population. *Nutrients* 2014; 6: 4961-4973.
  35. Hasima N, Aggarwal BB. Cancer-linked targets modulated by curcumin. *Int J Biochem Mol Biol* 2012; 3: 328.
  36. Russo M, Spagnuolo C, Tedesco I, et al. The flavonoid quercetin in disease prevention and therapy: facts and fancies. *Biochem Pharmacol* 2012; 83: 6-15.
  37. Cragg GM, Pezzuto JM. Natural products as a vital source for the discovery of cancer chemotherapeutic and chemopreventive agents. *Med Princ Pract* 2016; 25: 41-59.
  38. Ansari J, Inamdar N. The promise of traditional medicines. *Int J Pharmacol* 2010; 6: 808-812.
  39. Shukla S, Meeran SM. Epigenetics of cancer stem cells: Pathways and therapeutics. *Biochim Biophys Acta* 2014; 1840: 3494-3502.
  40. Popa V, Dumitru M, Volf I, et al. Lignin and polyphenols as allelochemicals. *Ind Crops Prod* 2008; 27: 144-148.
  41. Papadopoulou A, Frazier RA. Characterization of protein-polyphenol interactions. *Trends Food Sci Technol* 2004; 15: 186-190.
  42. Thomasset SC, Berry DP, Garcea G, et al. Dietary polyphenolic phytochemicals-promising cancer chemopreventive agents in humans? A review of their clinical properties. *Int J Cancer* 2007; 120: 451-458.
  43. Gonzalez-Vallinas M, Gonzalez-Castejon M, Rodriguez-Casado A, et al. Dietary phytochemicals in cancer prevention and therapy: a complementary approach with promising perspectives. *Nutr Rev* 2013; 71: 585-599.
  44. Bonfili L, Cecarini V, Amici M, et al. Natural polyphenols as proteasome modulators and their role as anti-cancer compounds. *FEBS J* 2008; 275: 5512-5526.
  45. Tait S, Salvati AL, Desideri N, et al. Antiviral activity of substituted homoisoflavonoids on enteroviruses. *Antiviral Res* 2006; 72: 252-255.
  46. Özçelik B, Orhan DD, Özgen S, et al. Antimicrobial activity of flavonoids against extended-spectrum  $\beta$ -lactamase (ES $\beta$ L)-producing *Klebsiella pneumoniae*. *Trop J Pharm Res* 2008; 7: 1151-1157.
  47. Hendrich AB. Flavonoid-membrane interactions: possible consequences for biological effects of some polyphenolic compounds. *Acta Pharmacol Sin* 2006; 27: 27-40.
  48. Arabbi PR, Genovese MI, Lajolo FM. Flavonoids in vegetable foods commonly consumed in Brazil and estimated ingestion by the Brazilian population. *J Agri Food Chem* 2004; 52: 1124-1131.
  49. Justesen U, Knuthsen P, Leth T. Quantitative analysis of flavonols, flavones, and flavanones in fruits, vegetables and beverages by high-performance liquid chromatography with photo-diode array and mass spectrometric detection. *J Chromatogr A* 1998; 799: 101-110.
  50. McCann SE, Ambrosone CB, Moysich KB, et al. Intakes
-

- 
- of selected nutrients, foods, and phytochemicals and prostate cancer risk in western New York. *Nutr Cancer* 2005; 53: 33-41.
51. Sakakibara H, Honda Y, Nakagawa S, et al. Simultaneous determination of all polyphenols in vegetables, fruits, and teas. *J Agri Food Chem* 2003; 51: 571-581.
52. Mattila P, Astola J, Kumpulainen J. Determination of flavonoids in plant material by HPLC with diode-array and electro-array detections. *J Agri Food Chem* 2000; 48: 5834-5841.
53. Khan N, Mukhtar H. Tea and health: studies in humans. *Curr Pharm Des* 2013; 19: 6141-6147.
54. Shah S, Stapleton P, Taylor P. The polyphenol (-)-epicatechin gallate disrupts the secretion of virulence-related proteins by *Staphylococcus aureus*. *Lett Appl Microbiol* 2008; 46: 181-185.
55. Nifli AP, Bosson-Kouamé A, Papadopoulou N, et al. Monomeric and oligomeric flavanols are agonists of membrane androgen receptors. *Exp Cell Res* 2005; 309: 329-339.
56. Kuzuhara T, Tanabe A, Sei Y, et al. Synergistic effects of multiple treatments, and both DNA and RNA direct bindings on, green tea catechins. *Mol Carcinog* 2007; 46: 640-645.
57. Singh BN, Shankar S, Srivastava RK. Green tea catechin, epigallocatechin-3-gallate (EGCG): mechanisms, perspectives and clinical applications. *Biochem Pharmacol* 2011; 82: 1807-1821.
58. Bhagwat S, Haytowitz DB, Holden JM. USDA database for the flavonoid content of selected foods, Release 3.1. US Department of Agriculture: Beltsville 2014; 11: 145-149.
59. Yang CS, Wang X, Lu G, et al. Cancer prevention by tea: animal studies, molecular mechanisms and human relevance. *Nat Rev Cancer* 2009; 9: 429.
60. Shi J, Liu F, Zhang W, et al. Epigallocatechin-3-gallate inhibits nicotine-induced migration and invasion by the suppression of angiogenesis and epithelial-mesenchymal transition in non-small cell lung cancer cells. *Oncol Rep* 2015; 33: 2972-2980.
61. Deng YT, Lin JK. EGCG inhibits the invasion of highly invasive CL1-5 lung cancer cells through suppressing MMP-2 expression via JNK signaling and induces G2/M arrest. *J Agri Food Chem* 2011; 59: 13318-13327.
62. Banerjee S. Inhibition of mackerel (*Scomber scombrus*) muscle lipoxygenase by green tea polyphenols. *Food Res Int* 2006; 39: 486-491.
63. Sen T, Dutta A, Amitava C. Epigallocatechin-3-gallate (EGCG) downregulates gelatinase-B (MMP-9) by involvement of FAK/ERK/NFkappaB and AP-1 in the human breast cancer cell line MDA-MB-231. *Anticancer Drugs* 2010; 21: 632-624.
64. Khan N, Mukhtar H. Modulation of signaling pathways in prostate cancer by green tea polyphenols. *Biochem Pharmacol* 2013; 85: 667-672.
65. Chuang JC, Yoo CB, Kwan JM, et al. Comparison of biological effects of non-nucleoside DNA methylation inhibitors versus 5-aza-2'-deoxycytidine. *Mol Cancer Ther* 2005; 4: 1515-1520.
66. Gilbert E, Liu D. Flavonoids influence epigenetic-modifying enzyme activity: structure-function relationships and the therapeutic potential for cancer. *Curr Med Chem* 2010; 17: 1756-1768.
67. Lin Z, Fischer J, Wicker L. Intermolecular binding of blueberry pectin-rich fractions and anthocyanin. *Food Chem* 2016; 194: 986-993.
68. Stoner GD, Wang LS, Casto BC. Laboratory and clinical studies of cancer chemoprevention by antioxidants in berries. *Carcinogenesis* 2008; 29: 1665-1674.
69. Bin Hafeez B, Asim M, Siddiqui IA, et al. Delphinidin, a dietary anthocyanidin in pigmented fruits and vegeta-
-

- 
- bles: a new weapon to blunt prostate cancer growth. *Cell Cycle* 2008; 7: 3320-3326.
70. Kausar H, Jeyabalan J, Aqil F, et al. Berry anthocyanidins synergistically suppress growth and invasive potential of human non-small-cell lung cancer cells. *Cancer Lett* 2012; 325: 54-62.
71. Liu W, Xu J, Wu S, et al. Selective anti-proliferation of HER2-positive breast cancer cells by anthocyanins identified by high-throughput screening. *PLoS One* 2013; 8: e81586.
72. Lee SJ, Hong S, Yoo SH, Kim GW. Cyanidin-3-O-sambubioside from *Acanthopanax sessiliflorus* fruit inhibits metastasis by downregulating MMP-9 in breast cancer cells MDA-MB-231. *Planta Med* 2013; 79: 1636-1640.
73. Murakami A, Ashida H, Terao J. Multitargeted cancer prevention by quercetin. *Cancer Lett* 2008; 269: 315-325.
74. Busch C, Burkard M, Leischner C, et al. Epigenetic activities of flavonoids in the prevention and treatment of cancer. *Clin Epigenetics* 2015; 7: 64-67.
75. Ferraresi R, Troiano L, Roat E, et al. Essential requirement of reduced glutathione (GSH) for the anti-oxidant effect of the flavonoid quercetin. *Free Radic Res* 2005; 39: 1249-1258.
76. Chen C, Zhou J, Ji C. Quercetin: a potential drug to reverse multidrug resistance. *Cell Mol Life Sci* 2010; 87: 333-338.
77. Yamazaki S, Miyoshi N, Kawabata K, et al. Quercetin-3-O-glucuronide inhibits noradrenaline-promoted invasion of MDA-MB-231 human breast cancer cells by blocking  $\beta$ 2-adrenergic signaling. *Arch Biochem Biophys* 2014; 557: 18-27.
78. Zamora-Ros R, Knaze V, Luján-Barroso L, et al. Estimated dietary intakes of flavonols, flavanones and flavones in the European Prospective Investigation into Cancer and Nutrition (EPIC) 24-hour dietary recall cohort. *Br J Nutr* 2011; 106: 1915-1925.
79. Murti Y, Mishra P. Synthesis and evaluation of flavanones as anticancer agents. *Indian J Pharm Sci* 2014; 76: 163.
80. Choi EJ. Hesperetin induced G1-phase cell cycle arrest in human breast cancer MCF-7 cells: involvement of CDK4 and p21. *Nutr Cancer* 2007; 59: 115-119.
81. Palit S, Kar S, Sharma G, et al. Hesperetin induces apoptosis in breast carcinoma by triggering accumulation of ROS and activation of ASK1/JNK pathway. *J Cell Physiol* 2015; 230: 1729-1739.
82. Alshatwi AA, Ramesh E, Periasamy V, et al. The apoptotic effect of hesperetin on human cervical cancer cells is mediated through cell cycle arrest, death receptor, and mitochondrial pathways. *Fundam Clin Pharmacol* 2013; 27: 581-592.
83. Sambantham S, Radha M, Paramasivam A, et al. Molecular mechanism underlying hesperetin-induced apoptosis by in silico analysis and in prostate cancer PC-3 cells. *Asian Pac J Cancer Prev* 2013; 14: 4347-4352.
84. Zhang J, Wu D, Song J, et al. Hesperetin induces the apoptosis of gastric cancer cells via activating mitochondrial pathway by increasing reactive oxygen species. *Dig Dis Sci* 2015; 60: 2985-2995.
85. Singh M, Kaur M, Silakari O. Flavones: An important scaffold for medicinal chemistry. *Eur J Med Chem* 2014; 84: 206-239.
86. Meyer H, Bolarinwa A, Wolfram G, et al. Bioavailability of apigenin from apiin-rich parsley in humans. *Ann Nutr Metab* 2006; 50: 167-172.
87. Das S, Das J, Samadder A, Paul A, et al. Strategic formulation of apigenin-loaded PLGA nanoparticles for intracellular trafficking, DNA targeting and improved therapeutic effects in skin melanoma in vitro. *Toxicol Lett*
-

- 
- 2013; 223: 124-138.
88. Pandey M, Kaur P, Shukla S, et al. Plant flavone apigenin inhibits HDAC and remodels chromatin to induce growth arrest and apoptosis in human prostate cancer cells: in vitro and in vivo study. *Mol Carcinog* 2012; 51: 952-962.
89. Duarte S, Arango D, Parihar A, et al. Apigenin protects endothelial cells from lipopolysaccharide (LPS)-induced inflammation by decreasing caspase-3 activation and modulating mitochondrial function. *Int J Mol Sci* 2013; 14: 17664-17679.
90. Arango D, Parihar A, Villamena FA, et al. Apigenin induces DNA damage through the PKC $\delta$ -dependent activation of ATM and H2AX causing down-regulation of genes involved in cell cycle control and DNA repair. *Biochem Pharmacol* 2012; 84: 1571-1580.
91. Kroonen J, Artesi M, Capraro V, et al. Casein kinase 2 inhibition modulates the DNA damage response but fails to radiosensitize malignant glioma cells. *Int J Oncol* 2012; 41: 776-782.
92. Gates MA, Vitonis AF, Tworoger SS, et al. Flavonoid intake and ovarian cancer risk in a population-based case-control study. *Int J Cancer* 2009; 124: 1918-1925.
93. Szkudelska K, Nogowski L. Genistein-a dietary compound inducing hormonal and metabolic changes. *J Steroid Biochem Mol Biol* 2007; 105: 37-45.
94. Wang H, Li Q, Chen H. Genistein affects histone modifications on Dickkopf-related protein 1 (DKK1) gene in SW480 human colon cancer cell line. *PLoS One* 2012; 7: e40955.
95. Kelly AM, Green CC. Binding affinities of hepatic nuclear estrogen receptors for genistein in channel catfish. *N Am J Aquac* 2006; 68: 160-167.
96. Tian T, Li J, Li B, et al. Genistein exhibits anti-cancer effects via down-regulating FoxM1 in H446 small-cell lung cancer cells. *Tumour Biol* 2014; 35: 4137-4145.
97. Fang MZ, Chen D, Sun Y, et al. Reversal of hypermethylation and reactivation of p16INK4a, RAR $\beta$ , and MGMT genes by genistein and other isoflavones from soy. *Clin Cancer Res* 2005; 11: 7033-7041.
98. Uckun F. Use of EGF genistein to prevent development of EGF-receptor expressing cancers. Parker Hughes Institute 2002; U.S. Patent Application 10/145,798
99. Qin J, Teng J, Zhu Z, et al. Genistein induces activation of the mitochondrial apoptosis pathway by inhibiting phosphorylation of Akt in colorectal cancer cells. *Pharm Biol* 2016; 54: 74-79.
100. Farah A, Donangelo CM. Phenolic compounds in coffee. *Braz J Plant Physiol* 2006; 18: 23-36.
101. Fiuza S, Gomes C, Teixeira L, et al. Phenolic acid derivatives with potential anticancer properties-a structure-activity relationship study. Part 1: Methyl, propyl and octyl esters of caffeic and gallic acids. *Bioorg Med Chem* 2004; 12: 3581-3589.
102. Spilioti E, Jaakkola M, Tolonen T, et al. Phenolic acid composition, antiatherogenic and anticancer potential of honeys derived from various regions in Greece. *PLoS One* 2014; 9(4): e94860.
103. Subramanian AP, Jaganathan SK, Mandal M, et al. Gallic acid induced apoptotic events in HCT-15 colon cancer cells. *World J Gastroenterol* 2016; 22(15): 3952-3961.
104. Ho HH, Chang CS, Ho WC, et al. Gallic acid inhibits gastric cancer cells metastasis and invasive growth via increased expression of RhoB, downregulation of AKT/small GTPase signals and inhibition of NF-kappaB activity. *Toxicol Appl Pharmacol* 2013; 266(1): 76-85.
105. Fabiani R, Rosignoli P, De Bartolomeo A, et al. The production of hydrogen peroxide is not a common mechanism by which olive oil phenols induce apoptosis
-

- 
- on HL60 cells. *Food Chem* 2011; 125(4): 1249-1255.
106. Zhao B and Hu M. Gallic acid reduce cell viability, proliferation, invasion and angiogenesis in human cervical cancer cells. *Oncol Lett* 2013; 6(6): 1749-1755.
107. Forester SC, Choy YY, Waterhouse AL, et al. The anthocyanin metabolites gallic acid, 3-O-methylgallic acid, and 2, 4, 6-trihydroxybenzaldehyde decrease human colon cancer cell viability by regulating pro-oncogenic signals. *Mol Carcinog* 2014; 53(6): 432-439.
108. Vallverdú-Queralt A, Verbaere A, Meudec E, et al. Straightforward method to quantify GSH, GSSG, GRP, and hydroxycinnamic acids in wines by UPLC-MRM-MS. *J Agri Food Chem* 2015; 63(1): 142-149.
109. Lin CL, Chen RF, Chen JYF, et al. Protective effect of caffeic acid on paclitaxel induced anti-proliferation and apoptosis of lung cancer cells involves NF- $\kappa$ B pathway. *Int J Mol Sci* 2012; 13(5): 6236-6245.
110. Prasad NR, Karthikeyan A, Karthikeyan S, et al. Inhibitory effect of caffeic acid on cancer cell proliferation by oxidative mechanism in human HT-1080 fibrosarcoma cell line. *Mol Cell Biochem* 2011; 349(1-2): 11-19.
111. Beejmohun V, Fliniaux O, Grand É, et al. Microwave-assisted extraction of the main phenolic compounds in flaxseed. *Phytochemical Analysis. Int J Plant Chem Biochem Tech* 2007; 18(4): 275-282.
112. Mota FL, Queimada AJ, Pinho SP, et al. Aqueous solubility of some natural phenolic compounds. *Ind Eng Chem Res* 2008; 47(15): 5182-5189.
113. Karthikeyan S, Kanimozhi G, Prasad NR, et al. Radiosensitizing effect of ferulic acid on human cervical carcinoma cells in vitro. *Toxicol In Vitro* 2011; 25(7): 1366-1375.
114. Eroğlu C, Seçme M, Bağcı G, et al. Assessment of the anticancer mechanism of ferulic acid via cell cycle and apoptotic pathways in human prostate cancer cell lines. *Tumor Biol* 2015; 36(12): 9437-9446.
115. Milder IE, Arts IC, van de Putte B, et al. Lignan contents of Dutch plant foods: a database including larciresinol, pinoresinol, secoisolariciresinol and matairesinol. *Br J Nutr* 2005; 93(3): 393-402.
116. Webb AL and McCullough ML. Dietary lignans: potential role in cancer prevention. *Nutr Cancer* 2005; 51(2): 117-131.
117. Linseisen J, Piller R, Hermann S, et al. Dietary phytoestrogen intake and premenopausal breast cancer risk in a German case-control study. *Int J Cancer* 2004; 110(2): 284-290.
118. Deng P, Wang C, Chen L, et al. Sesamin induces cell cycle arrest and apoptosis through the inhibition of signal transducer and activator of transcription 3 signaling in human hepatocellular carcinoma cell line HepG2. *Biol Pharm Bull* 2013; 36(10): 1540-1548.
119. Xu P, Cai F, Liu X, et al. Sesamin inhibits lipopolysaccharide-induced proliferation and invasion through the p38-MAPK and NF- $\kappa$ B signaling pathways in prostate cancer cells. *Oncol Rep* 2015; 33(6): 3117-3123.
120. King RE, Bomser JA and Min DB. Bioactivity of resveratrol. *Compr Rev Food Sci Food Saf* 2006; 5: 65-70.
121. Anekonda TS. Resveratrol-a boon for treating Alzheimer's disease? *Brain Res Rev* 2006; 52(2): 316-326.
122. Kraft TE, Parisotto D, Schempp C, et al. Fighting cancer with red wine? Molecular mechanisms of resveratrol. *Crit Rev Food Sci Nutr* 2009; 49(9): 782-799.
123. Athar M, Back JH, Tang X, et al. Resveratrol: a review of preclinical studies for human cancer prevention. *Toxicol App Pharmacol* 2007; 224(3): 274-283.
124. Yang Q, Wang B, Zang W, et al. Resveratrol inhibits the growth of gastric cancer by inducing G1 phase arrest and senescence in a Sirt1-dependent manner. *PLoS One* 2013; 8(11): e70627.
125. Aluyen JK, Ton QN, Tran T, et al. Resveratrol: potential as anticancer agent. *J Diet Suppl* 2012; 9(1): 45-56.
-

- 
126. Scarlatti F, Maffei R, Beau I, et al. Role of non-canonical Beclin 1-independent autophagy in cell death induced by resveratrol in human breast cancer cells. *Cell Death Differ* 2008; 15(8): 1318-1329.
127. Gao X, Xu YX, Divine G, et al. Disparate in vitro and in vivo antileukemic effects of resveratrol, a natural polyphenolic compound found in grapes. *J Nutr* 2002; 132(7): 2076-2081.
128. Wenzel E, Soldo T, Erbersdobler H, et al. Bioactivity and metabolism of trans-resveratrol orally administered to Wistar rats. *Mol Nutr Food Res* 2005; 49(5): 482-494.
129. Madan E, Prasad S, Roy P, et al. Regulation of apoptosis by resveratrol through JAK/STAT and mitochondria mediated pathway in human epidermoid carcinoma A431 cells. *Biochem Biophys Res Commun* 2008; 377(4): 1232-1237.
130. Li H, Wu WKK, Zheng Z, et al. 3, 3', 4, 5, 5'-penta-hydroxy-trans-stilbene, a resveratrol derivative, induces apoptosis in colorectal carcinoma cells via oxidative stress. *Eur J Pharmacol* 2010; 637(1-3): 55-61.
131. Brizuela L, Dayon A, Doumerc N, et al. The sphingosine kinase-1 survival pathway is a molecular target for the tumor-suppressive tea and wine polyphenols in prostate cancer. *FASEB J* 2010; 24(10): 3882-3894.
132. Cai Y, Luo Q, Sun M, et al. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 2004; 74(17): 2157-2184.
133. Ghosh R, Nadiminty N, Fitzpatrick JE, et al. Eugenol causes melanoma growth suppression through inhibition of E2F1 transcriptional activity. *J Biol Chem* 2005; 280(7): 5812-5819.
134. Sun J, Yao J, Huang S, et al. Antioxidant activity of polyphenol and anthocyanin extracts from fruits of *Kadsura coccinea* (Lem.) AC Smith. *Food Chem* 2009; 117(2): 276-281.
135. Tu SH, Ku CY, Ho CT, et al. Tea polyphenol (-)-epigallocatechin-3-gallate inhibits nicotine-and estrogen-induced  $\alpha$ 9-nicotinic acetylcholine receptor upregulation in human breast cancer cells. *Mol Nutr Food Res* 2011; 55(3): 455-466.
136. Ronnekleiv-Kelly SM, Nukaya M, Díaz-Díaz CJ, et al. Aryl hydrocarbon receptor-dependent apoptotic cell death induced by the flavonoid chrysin in human colorectal cancer cells. *Cancer Lett* 2016; 370(1): 91-99.
137. Duo J, Ying GG, Wang GW, et al. Quercetin inhibits human breast cancer cell proliferation and induces apoptosis via Bcl-2 and Bax regulation. *Mol Med Rep* 2012; 5(6): 1453-1456.
138. Magee PJ, Allsopp P, Samaletdin A, et al. Daidzein, R-(+) equol and S(-) equol inhibit the invasion of MDA-MB-231 breast cancer cells potentially via the down-regulation of matrix metalloproteinase-2. *Eur J Nutr* 2014; 53(1): 345-350.
139. Rigalli JP, Tocchetti GN, Arana MR, et al. The phytoestrogen genistein enhances multidrug resistance in breast cancer cell lines by translational regulation of ABC transporters. *Cancer Lett* 2016; 376(1): 165-172.
140. Mahmoud AM, Zhu T, Parray A, et al. Differential effects of genistein on prostate cancer cells depend on mutational status of the androgen receptor. *PLoS One* 2013; 8(10): e78479.
141. Chen HS, Bai MH, Zhang T, et al. Ellagic acid induces cell cycle arrest and apoptosis through TGF-beta/Smad3 signaling pathway in human breast cancer MCF-7 cells. *Int J Oncol* 2015; 46(4): 1730-1738.
142. Yong WK, Ho YF and Malek SNA. Xanthohumol induces apoptosis and S phase cell cycle arrest in A549 non-small cell lung cancer cells. *Pharmacogn Mag* 2015; 11(2): S275-S283.
143. Jin CY, Park C, Hwang HJ, et al. Naringenin up-regu-
-

- 
- lates the expression of death receptor 5 and enhances TRAIL-induced apoptosis in human lung cancer A549 cells. *Mol Nutr Food Res* 2011; 55(2): 300-309.
144. Hong Z, Cao X, Li N, et al. Luteolin is effective in the non-small cell lung cancer model with L858R/T790M EGF receptor mutation and erlotinib resistance. *Br J Pharmacol* 2014; 171(11): 2842-2853.
145. Siddiqui IA, Asim M, Hafeez BB, et al. Green tea polyphenol EGCG blunts androgen receptor function in prostate cancer. *FASEB J* 2011; 25(4): 1198-1207.
146. Shukla S, Bhaskaran N, Babcook MA, et al. Apigenin inhibits prostate cancer progression in TRAMP mice via targeting PI3K/Akt/FoxO pathway. *Carcinogenesis* 2013; 35(2): 452-460.
147. Pitchakarn P, Chewonarin T, Ogawa K, et al. Ellagic acid inhibits migration and invasion by prostate cancer cell lines. *Asian Pac J Cancer Prev* 2013; 14(5): 2859-2863.
148. Ganapathy S, Chen Q, Singh KP, et al. Resveratrol enhances antitumor activity of TRAIL in prostate cancer xenografts through activation of FOXO transcription factor. *PloS One* 2010; 5(12): e15627.
149. Selvaraj S, Sun Y, Sukumaran P, et al. Resveratrol activates autophagic cell death in prostate cancer cells via downregulation of STIM1 and the mTOR pathway. *Mol Carcinog* 2016; 55(5): 818-831.
150. Yun JM, Afaq F, Khan N, et al. Delphinidin, an anthocyanidin in pigmented fruits and vegetables, induces apoptosis and cell cycle arrest in human colon cancer HCT116 cells. *Mol Carcinog* 2009; 48(3): 260-270.
151. Thakur VS, Deb G, Babcook MA, et al. Plant phytochemicals as epigenetic modulators: role in cancer chemoprevention. *AAPS J* 2014; 16(1): 151-163.
152. Song HM, Park GH, Eo HJ, et al. Naringenin-Mediated ATF3 Expression Contributes to Apoptosis in Human Colon Cancer. *Biomol Ther* 2016; 24(2): 140-146.
153. Chunhua L, Donglan L, Xiuqiong F, et al. Apigenin up-regulates transgelin and inhibits invasion and migration of colorectal cancer through decreased phosphorylation of AKT. *J Nutr Biochem* 2013; 24(10): 1766-1775.
154. Lim DY, Cho HJ, Kim J, et al. Luteolin decreases IGF-II production and downregulates insulin-like growth factor-I receptor signaling in HT-29 human colon cancer cells. *BMC Gastroenterol* 2012; 12: 9-14.
155. Lee HS, Cho HJ, Yu R, et al. Mechanisms underlying apoptosis-inducing effects of Kaempferol in HT-29 human colon cancer cells. *Int J Mol Sci* 2014; 15(2): 2722-2737.
156. Yeh CB, Hsieh MJ, Lin CW, et al. The antimetastatic effects of resveratrol on hepatocellular carcinoma through the downregulation of a metastasis-associated protease by SP-1 modulation. *PloS One* 2013; 8(3): e56661.
157. Hussain A, Harish G, Prabhu GA, et al. Inhibitory effect of genistein on the invasive potential of human cervical cancer cells via modulation of matrix metalloproteinase-9 and tissue inhibitors of matrix metalloproteinase-1 expression. *Cancer Epidemiol* 2012; 36(6): e387-e393.
158. Bishayee K, Ghosh S, Mukherjee A, et al. Quercetin induces cytochrome-c release and ROS accumulation to promote apoptosis and arrest the cell cycle in G2/M, in cervical carcinoma: signal cascade and drug-DNA interaction. *Cell Prolif* 2013; 46(2): 153-163.
159. García-Zepeda SP, García-Villa E, Díaz-Chávez J, et al. Resveratrol induces cell death in cervical cancer cells through apoptosis and autophagy. *Eur J Cancer Prev* 2013; 22(6): 577-584.
160. Zhao X, Jiang K, Liang B, et al. Anticancer effect of xanthohumol induces growth inhibition and apoptosis of human liver cancer through NF- $\kappa$ B/p53-apoptosis signaling pathway. *Oncol Rep* 2016; 35(2): 669-675.
161. Park HJ, Jeon YK, You DH, et al. Daidzein causes cy-
-

- 
- tochrome c-mediated apoptosis via the Bcl-2 family in human hepatic cancer cells. *Food Chem Toxicol* 2013; 60: 542-549.
162. Srigopalram S, Jayraaj I, Kaleeswaran B, et al. Elagic acid normalizes mitochondrial outer membrane permeabilization and attenuates inflammation-mediated cell proliferation in experimental liver cancer. *App Biochem Biotech* 2014; 173(8): 2254-2266.
163. Rajasekaran D, Elavarasan J, Sivalingam M, et al. Resveratrol interferes with N-nitrosodiethylamine-induced hepatocellular carcinoma at early and advanced stages in male Wistar rats. *Mol Med Rep* 2011; 4: 1211-1217.
164. Onoda C, Kuribayashi K, Nirasawa S, et al. (-)-Epigallocatechin-3-gallate induces apoptosis in gastric cancer cell lines by down-regulating survivin expression. *Int J Oncol* 2011; 38(5): 1403-1408.
165. Chen M, Rao Y, Zheng Y, et al. Association between soy isoflavone intake and breast cancer risk for pre- and post-menopausal women: A meta-analysis of epidemiological studies. *PLoS ONE* 2014; 9(2): e89288-e89292.
166. Fresco P, Borges F, Diniz C, et al. New insights on the anticancer properties of dietary polyphenols. *Med Res Rev* 2006; 26(6): 747-766.
167. Han X, Shen T and Lou H. Dietary polyphenols and their biological significance. *Int J Mol Sci* 2007; 8(9): 950-988
168. Luo Y, Wang SX, Zhou ZQ, et al. Apoptotic effect of genistein on human colon cancer cells via inhibiting the nuclear factor-kappa B (NF- $\kappa$ B) pathway. *Tumor Biol* 2014; 35(1): 11483-11488.
169. Kwon KH, Barve A, Yu S, et al. Cancer chemoprevention by phytochemicals: potential molecular targets, biomarkers and animal models 1. *Acta Pharmacol Sin* 2007; 28(9): 1409-1421.
-



